

Immunization Timing

Age 2 months DTaP HepB* Hib PCV (Pneumo) Polio (IPV)	Age 4 months DTaP HepB* Hib PCV (Pneumo) Polio (IPV)	Age 6 months DTaP HepB* Hib[†] PCV (Pneumo) Polio (IPV)	Age 12 months MMR Varicella Hib PCV (Pneumo)	Age 15 months DTaP	Age 2 years HepA
	Interval from previous dose 1-2 months 1-2 months 1-2 months 1-2 months 1-2 months	Interval from previous dose 1-2 months 2-12 months and at least 4 months after first dose 1-2 months 1-2 months 2-14 months	Interval from previous dose On or after first birthday On or after first birthday 2-8 months and on or after first birthday 6-8 weeks and on or after first birthday	Interval from previous dose 6-12 months	Interval from previous dose 2nd dose 6-12 months after first

Every Fall: Flu Vaccine

- All children 6–23 months of age.
- All children 6 months and older with asthma, diabetes, or other chronic high-risk condition.
- Others in the household of both of the above.

Note: Children 6 months through 8 years old need 2 doses, one month apart, in their first year of flu shots.


Before Kindergarten

Polio, DTaP, and MMR #2.
Also, unless already given, HepB and Varicella.

This is a suggested schedule. For alternatives and details, consult the latest "Recommended Childhood Immunization Schedule, United States."

* Hep B series also can be started at birth, the second dose given at 1–2 months and the third dose at 6–12 months. If using Comvax[®] (combined Hib and HepB vaccine), the third dose should be given at 12 months, not 6 months.

† This dose can be skipped if Pedvax Hib[®] and/or Comvax[®] (combined Hib and HepB vaccine) are used exclusively for the infant doses.



CALIFORNIA KIDS
Love them. Immunize them.